

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 335 250 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
13.08.2003 Bulletin 2003/33

(51) Int Cl.7: **G03G 5/06**, C07D 333/48,
C07C 251/84, C07C 317/34,
C07D 249/18, C07D 231/38,
C07D 257/06, C07D 207/34

(21) Application number: 03250767.5

(22) Date of filing: 06.02.2003

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT SE SI SK TR
Designated Extension States:
AL LT LV MK RO

(30) Priority: 08.02.2002 US 355018 P
08.02.2002 US 355047 P
08.02.2002 US 355060 P
08.02.2002 US 355228 P
08.02.2002 US 355019 P
08.02.2002 US 355080 P
08.02.2002 US 355079 P
08.02.2002 US 355073 P
08.02.2002 US 355066 P

(71) Applicant: **SAMSUNG ELECTRONICS CO., LTD.**
Suwon-City, Kyungki-do (KR)


(72) Inventors:
• Law, Kam W.
Woodbury, MN 55125 (US)
• Jubran, Nusrallah
St. Paul, MN 55119 (US)
• Tokarski, Zbigniew
Woodbury, MN 55125 (US)
• Katritzky, Alan R.
Gainesville, FL 32601 (US)
• Jain, Ritu
Gainesville, FL 32601 (US)
• Malmait, Rexiat
Cincinnati, OH 45249 (US)

(74) Representative: **Moy, David et al**
Appleyard Lees,
15 Clare Road
Halifax HX1 2HY (GB)

(54) **Electrophotographic organophotoreceptors with novel charge transport materials**

(57) This invention relates to a novel organophotoreceptor comprising:

(a) at least one charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolylhydrazone groups, d) at least two fluorenone sulfolanylhydrazone groups, e) at least two fluorenone pyrazolylhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolylhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups. Some of these fluorenones may be represented by the formula



where n is an integer between 2 and 6, inclusive; R₁ is hydrogen, an alkyl group, or an aryl group (e.g., phenyl, naphthyl, stilbenyl, benzyl, or tolyl group); R₂ is an alkylsulfonylphenyl or one of its derivatives; X is a

BEST AVAILABLE COPY

EP 1 335 250 A1

linking group having the formula $-(CH_2)_m-$, branched or linear. The compounds may form electrostatic imaging systems in combination with

(b) a charge generating compound; and

(c) an electrically conductive substrate.

Description

[0001] This invention relates to organophotoreceptors suitable for use in electrophotography and, more specifically, to flexible organophotoreceptors having novel charge transport materials. The generic class of charge transport materials are fluorenone hydrazones and have a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolylhydrazone groups, d) at least two fluorenone sulfolanylhydrazone groups, e) at least two fluorenone pyrazolylhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolylhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups.

[0002] In electrophotography, an organophotoreceptor in the form of a plate, a flexible belt, a disk, a rigid drum, or a sheet around a rigid or compliant drum having an electrically insulating photoconductive element on an electrically conductive substrate is imaged by first uniformly electrostatically charging the surface of the photoconductive layer, and then exposing the charged surface to a pattern of light. The light exposure selectively dissipates the charge in the illuminated areas, thereby forming a pattern of charged and uncharged areas (referred to as latent image). A liquid or solid toner is then provided in the vicinity of the latent image, and the toner particles deposit in either the charged or uncharged areas to create a toned image on the surface of the photoconductive layer. The resulting visible toner image can be transferred to a suitable receiving surface such as paper, or the photoconductive layer can operate as a permanent receptor for the image. The imaging process can be repeated many times.

[0003] Both single layer and multilayer photoconductive elements have been used. In the single layer embodiment, a charge transport material and charge generating material are combined with a polymeric binder and then deposited on an electrically conductive substrate. In the multilayer embodiment, the charge transport material and charge generating material are in the form of separate layers, each of which can optionally be combined with a polymeric binder and deposited on the electrically conductive substrate. Two arrangements are possible. In one arrangement (the "dual layer" arrangement), the charge generating layer is deposited on the electrically conductive substrate and the charge transport layer is deposited on top of the charge generating layer. In an alternate arrangement (the "inverted dual layer" arrangement), the order of the charge transport layer and charge generating layer is reversed.

[0004] In both the single layer and multilayer photoconductive elements, the purpose of the charge generating material is to generate charge carriers (i.e., holes or electrons) upon exposure to light. The purpose of the charge transport material is to accept these charge carriers and transport them through the charge transport layer in order to discharge a surface charge on the photoconductive element. The charge transport material can be a charge transport compound, an electron transport compound, or a combination of both. When a charge transport compound is used, the charge transport compound accepts the hole carriers and transports them through the layer containing the charge transport compound. When an electron transport compound is used, the electron transport compound accepts the electron carriers and transports them through the layer containing the electron transport compound.

[0005] To produce high quality images, it is desirable to maximize the amount of charge which the charge transport material, such as electron transport compound, can accept (indicated by a parameter known as the acceptance voltage or " V_{acc} "), and to minimize retention of that charge upon discharge (indicated by a parameter known as the discharge voltage or " V_{dis} ").

[0006] There are many charge transport materials available for electrophotography. The most common charge transport materials are pyrazoline derivatives, fluorene derivatives, oxadiazole derivatives, stilbene derivatives, hydrazone derivatives, carbazole hydrazone derivatives, triphenylamine derivatives, julolidine hydrazone derivatives, polyvinyl carbazole, polyvinyl pyrene, or polyacenaphthylene. However, each of the above charge transport materials suffers some disadvantages. There is always a need for novel charge transport materials to meet the various requirements of electrophotography applications.


[0007] According to the present invention there is provided an organophotoreceptor, an electrophotographic imaging apparatus, an electrophotographic imaging process, and a charge transfer material as set forth in the appended claims. Preferred features of the invention will be apparent from the dependent claims, and the description which follows.

[0008] In a first aspect, the invention features an organophotoreceptor comprising:

- a) at least one charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolylhydrazone groups, d) at least two fluorenone sulfolanylhydrazone groups, e) at least two fluorenone pyrazolylhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolylhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups;
- (b) a charge generating compound; and

(c) an electrically conductive substrate.

[0009] The charge transport material may be represented by the following formula



where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group (e.g., phenyl, naphthyl, stilbenyl, benzyl, or tolyl group, see U.S. Patent Nos. 6,469,123; 6,423,811; and 6,313,185 for a complete description and structure of this group, which patents are incorporated herein by reference for the disclosure of that group);

R₂ is an alkylsulfonylphenyl group, a pyrrolyl group, a benzotriazolyl group, a sulfolanyl group, a pyrazolyl group, a naphthyl group, a tetrazolyl group, a stilbenyl group, a (9H-fluoren-9-ylidene)benzyl group, or one of their derivatives having substituents thereon that do not alter the underlying elemental content or bond structure (e.g., changing double bonds to single bonds and single bonds to double bonds) of the recited group;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group (e.g., a cyclotetrasiloxyl group), a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group.

[0010] The charge transport material may or may not be symmetrical. Thus, for example, a linking group X for any given "arm" of the compound may be the same or different from the linking groups in other "arms" of the compound. Similarly, R₁ and R₂ groups for any given "arm" of the compound may be the same or different from R₁ and R₂ groups in any other arm. In addition, the above-described formula for the charge transport material is intended to cover isomers.

The organophotoreceptor may be provided in the form of a plate, a flexible belt, a disk, a rigid drum, or a sheet around a rigid or compliant drum. In one embodiment, the organophotoreceptor includes: (a) a charge transport layer comprising the charge transport material and a polymeric binder; (b) a charge generating layer comprising the charge generating compound and a polymeric binder; and (c) the electrically conductive substrate. The charge transport layer may be intermediate between the charge generating layer and the electrically conductive substrate. Alternatively, the charge generating layer may be intermediate between the charge transport layer and the electrically conductive substrate.


[0011] In various aspects, the invention features an electrophotographic imaging apparatus that includes (a) a plurality of support rollers; and (b) the above-described organophotoreceptor in the form of a flexible belt threaded around the support rollers. The apparatus preferably further includes a liquid toner dispenser. In another aspect, the invention features an electrophotographic imaging process that includes (a) applying an electrical charge to a surface of the above-described organophotoreceptor; (b) imagewise exposing the surface of the organophotoreceptor to radiation to dissipate charge in selected areas and thereby form a pattern of charged and uncharged areas on the surface; (c) contacting the surface with a liquid toner that includes a dispersion of colorant particles in an organic liquid to create a toned image; and (d) transferring the toned image to a substrate. In still another aspect, the invention features a novel charge transport material having Formula 1 above.

[0012] These organophotoreceptors can be used successfully with liquid toners to produce high quality images. The high quality of the images can be maintained after repeated cycling. Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

[0013] The invention features organophotoreceptors that include charge transport materials having the formula set forth as Formula 1 above. In addition, various subgeneric formulae may represent compounds of a more limited scope that are useful in the practice of the present invention. In addition to the scope of:

- a) at least one charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolylhydrazone groups, d) at least two fluorenone sulfolanylhydrazone groups, e) at least two fluorenone pyrazolylhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolylhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups;
- (b) a charge generating compound; and
- (c) an electrically conductive substrate,

the charge transport material may be represented by the following formula



where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group (e.g., phenyl, naphthyl, stilbenyl, benzyl, or tolyl group, see U.S. Patent Nos. 6,469,123; 6,423,811; and 6,313,185 for a complete description and structure of this group, which patents are incorporated herein by reference for the disclosure of that group);





R_2 is an alkylsulfonylphenyl group, a pyrrolyl group, a benzotriazolyl group, a sulfolanyl group, a pyrazolyl group, a naphthyl group, a tetrazolyl group, a stilbenyl group, a (9H-fluoren-9-ylidene)benzyl group, or one of their derivatives;




X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and




Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group (e.g., a cyclotetrasiloxyl group), a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.




[0014] Preferably R_1 and R_2 are, independently naphthyl or one of its derivatives.




[0015] Narrower descriptions of subgroups of compounds within the generic scope of the invention include, by way of non-limiting examples,















[0016] The electrically conductive substrate may be flexible, for example in the form of a flexible web or a belt, or inflexible, for example in the form of a drum. Typically, a flexible electrically conductive substrate comprises of an insulated substrate and a thin layer of electrically conductive materials. The insulated substrate may be paper or a film forming polymer such as polyethylene terephthalate, polyimide, polysulfone, polyethylene naphthalate, polypropylene, nylon, polyester, polycarbonate, polyvinyl fluoride, polystyrene and the like. Specific examples of supporting substrates included polyethersulfone (Stabar S-100, available from ICI), polyvinyl fluoride (Tedlar, available from E.I. DuPont de Nemours & Company), polybisphenol-A polycarbonate (Makrofol, available from Mobay Chemical Company) and amorphous polyethylene terephthalate (Melinar, available from ICI Americas, Inc.). The electrically conductive materials may be graphite, dispersed carbon black, iodide, conductive polymers such as polypyrroles and Calgon Conductive Polymer 261 (commercially available from Calgon Corporation, Inc., Pittsburgh, Pa.), metals such as aluminum, titanium, chromium, brass, gold, copper, palladium, nickel, or stainless steel, or metal oxide such as tin oxide or indium oxide. Preferably, the electrically conductive material is aluminum. Typically, the photoconductor substrate will have a thickness adequate to provide the required mechanical stability. For example, flexible web substrates generally have a thickness from about 0.01 to about 1 mm, while drum substrates generally have a thickness of from about 0.5 mm to about 2 mm.

[0017] The charge generating compound is a material which is capable of absorbing light to generate charge carriers, such as a dyestuff or pigment. Examples of suitable charge generating compounds include metal-free phthalocyanines, metal phthalocyanines such as titanium phthalocyanine, copper phthalocyanine, oxytitanium phthalocyanine, hydroxygallium phthalocyanine, squarylium dyes and pigments, hydroxy-substituted squarylium pigments, perylimides, polynuclear quinones available from Allied Chemical Corporation under the tradename Indofast™ Double Scarlet, Indofast™ Violet Lake B, Indofast™ Brilliant Scarlet and Indofast™ Orange, quinacridones available from DuPont under the tradename Monastral™ Red, Monastral™ Violet and Monastral™ Red Y, naphthalene 1,4,5,8-tetracarboxylic acid

derived pigments including the perinones, tetrabenzoporphyrins and tetranaphthaloporphyrins, indigo- and thioindigo dyes, benzothioxanthene-derivatives, perylene 3,4,9,10-tetracarboxylic acid derived pigments, polyazo-pigments including bisazo-, trisazo- and tetrakisazo-pigments, polymethine dyes, dyes containing quinazoline groups, tertiary amines, amorphous selenium, selenium alloys such as selenium-tellurium, selenium-tellurium-arsenic and selenium-arsenic, cadmium sulfoselenide, cadmiumselenide, cadmium sulfide, and mixtures thereof. Preferably, the charge generating compound is oxytitanium phthalocyanine, hydroxygallium phthalocyanine or a combination thereof.

[0018] Preferably, the charge generation layer comprises a binder in an amount of from about 10 to about 90 weight percent and more preferably in an amount of from about 20 to about 75 weight percent, based on the weight of the charge generation layer.

[0019] The binder is capable of dispersing or dissolving the charge transport material (in the case of the charge transport layer) and the charge generating compound (in the case of the charge generating layer). Examples of suitable binders for both the charge generating layer and charge transport layer include polystyrene-co-butadiene, modified acrylic polymers, polyvinyl acetate, styrene-alkyd resins, soya-alkyl resins, polyvinylchloride, polyvinylidene chloride, polyacrylonitrile, polycarbonates, polyacrylic acid, polyacrylates, polymethacrylates, styrene polymers, polyvinyl butyral, alkyd resins, polyamides, polyurethanes, polyesters, polysulfones, polyethers, polyketones, phenoxy resins, epoxy resins, silicone resins, polysiloxanes, poly(hydroxyether) resins, polyhydroxystyrene resins, novolak resins, resol resins, poly(phenylglycidyl ether)-co-dicyclopentadiene, copolymers of monomers used in the above-mentioned polymers, and combinations thereof. Polycarbonate binders and polyvinyl butyral are particularly preferred. Examples of suitable polycarbonate binders include polycarbonate A which is derived from bisphenol-A, polycarbonate Z, which is derived from cyclohexylidene bisphenol, polycarbonate C, which is derived from methylbisphenol A, and polyester carbonates. Examples of suitable polyvinyl butyral are BX-1 and BX-5 from Sekisui Chemical Co. Ltd., Japan.


[0020] Optionally, the organophotoreceptor may include a second charge transport material as known in the art. The charge transport material can be a charge transport compound, an electron transport compound, or a combination of both. When a charge transport compound is used, the charge transport compound accepts the hole carriers and transports them through the layer containing the charge transport compound. When an electron transport compound is used, the electron transport compound accepts the electron carriers and transports them through the layer containing the electron transport compound.

[0021] Suitable charge transport compounds include, but are not limited to, pyrazoline derivatives, fluorene derivatives, oxadiazole derivatives, stilbene derivatives, hydrazone derivatives, carbazole hydrazone derivatives, triaryl amines, polyvinylcarbazole, polyvinylpyrene, polyacenaphthylene, or multi-hydrazone compounds comprising at least two hydrazone groups and at least two groups selected from the group consisting of triphenylamine and heterocycles such as carbazole, julolidine, phenothiazine, phenazine, phenoxazine, phenoxathiin, thiazole, oxazole, isoxazole, dibenzo(1,4)dioxine, thianthrene, imidazole, benzothiazole, benzotriazole, benzoxazole, benzimidazole, quinoline, isoquinoline, quinoxaline, indole, indazole, pyrrole, purine, pyridine, pyridazine, pyrimidine, pyrazine, triazole, oxadiazole, tetrazole, thiadiazole, benzisoxazole, benzisothiazole, dibenzofuran, dibenzothiophene, thiophene, thianaphthene, quinazoline, or cinnoline.

[0022] Non-limiting examples of suitable electron transport compound include bromoanil, tetracyanoethylene, tetracyanoquinodimethane, 2,4,7-trinitro-9-fluorenone, 2,4,5,7-tetranitro-9-fluorenone, 2,4,5,7-tetranitroxanthone, 2,4,8-trinitrothioxanthone, 2,6,8-trinitro-indeno[1,2-b]thiophene-4-one, and 1,3,7-trinitro-dibenzothiophene-5,5-dioxide, (2,3-diphenyl-1-indenylidene)malononitrile, 4H-thiopyran-1,1-dioxide and its derivatives such as 4-dicyanomethylene-2,6-diphenyl-4H-thiopyran-1,1-dioxide, 4-dicyanomethylene-2,6-di-m-tolyl-4H-thiopyran-1,1-dioxide, and unsymmetrically substituted 2,6-diaryl-4H-thiopyran-1,1-dioxide such as 4H-1,1-dioxo-2-(p-isopropylphenyl)-6-phenyl-4-(dicyanomethylidene)thiopyran and 4H-1,1-dioxo-2-(p-isopropylphenyl)-6-(2-thienyl)-4-(dicyanomethylidene)thiopyran, derivatives of phospho-2,5-cyclohexadiene, alkoxycarbonyl-9-fluorenylidene)malononitrile derivatives such as (4-n-butoxycarbonyl-9-fluorenylidene)malononitrile, (4-phenethoxycarbonyl-9-fluorenylidene)malononitrile, (4-carbitoxy-9-fluorenylidene)malononitrile, and diethyl(4-n-butoxycarbonyl-2,7-dinitro-9-fluorenylidene)-malonate, anthraquinodimethane derivatives such as 11,11,12,12-tetracyano-2-alkylanthraquinodimethane and 11,11-dicyano-12,12-bis(ethoxycarbonyl)anthraquinodimethane, anthrone derivatives such as 1-chloro-10-[bis(ethoxycarbonyl)methylene]anthrone, 1,8-dichloro-10-[bis(ethoxycarbonyl)methylene]anthrone, 1,8-dihydroxy-10-[bis(ethoxycarbonyl)methylene]anthrone, and 1-cyano-10-[bis(ethoxycarbonyl)methylene]anthrone, 7-nitro-2-aza-9-fluorenylidene-malononitrile, diphenoquinone derivatives, benzoquinone derivatives, naphtoquinone derivatives, quinine derivatives, tetracyanoethylenecyanoethylene, 2,4,8-trinitrothioxanthone, dinitrobenzene derivatives, dinitroanthracene derivatives, dinitroacridine derivatives, nitroanthraquinone derivatives, dinitroanthraquinone derivatives, succinic anhydride, maleic anhydride, dibromo maleic anhydride, pyrene derivatives, carbazole derivatives, hydrazone derivatives, N,N-dialkylaniline derivatives, diphenylamine derivatives, triphenylamine derivatives, triphenylmethane derivatives, tetracyanoquinonedimethane, 2,4,5,7-tetranitro-9-fluorenone, 5,2,4,7-trinitro-9-dicyanomethylenefluorenone, 2,4,5,7-tetranitroxanthone derivatives, and 2,4,8-trinitrothioxanthone derivatives.

[0023] Optionally, the photoconductive layer of this invention may contain a light stabilizer. Non-limiting examples of

suitable light stabilizer include hindered trialkylamines such as Tinuvin 292 (from Ciba Specialty Chemicals, Terrytown, NY), hindered alkoxydialkylamines such as Tinuvin 123 (from Ciba Specialty Chemicals), benzotriazoles such as Tinuvin 928 (from Ciba Specialty Chemicals), benzophenones, nickel compounds such as Arbestab (from Robinson Brothers Ltd, West Midlands, Great Britain), salicylates, cyanocinnamates, benzylidene malonates, benzoates, oxanilides, polymeric sterically hindered amines such as Luchern (from atchem North America, Buffalo, NY). Preferably, the light stabilizer is selected from the group consisting of hindered trialkylamines having the following formula:



where $R_1, R_2, R_3, R_4, R_6, R_7, R_8, R_{10}, R_{11}, R_{12}, R_{13}, R_{15}, R_{16}$ are, independently, hydrogen, alkyl group, or ester, or ether group; and R_5, R_9 , and R_{14} are, independently, alkyl group; and X is a linking group selected from the group consisting of $-O-CO-(CH_2)_m-CO-O-$ where m is between 2 to 20.

[0024] The light stabilizer in the photoconductive layer is in an amount of from about 0.5 to about 25 weight percent and more preferably in an amount of from about 1 to about 10 weight percent, based on the weight of the photoconductive layer.

[0025] Conveniently, the photoconductive layer may be formed by dispersing or dissolving the components such as a charge generating compound, a charge transport compound, a light stabilizer, an electron transport compound, and a polymeric binder in organic solvent, coating the dispersion and/or solution on the respective underlying layer and drying the coating. Preferably, the components are dispersed by high shear homogenization, ball-milling, attritor milling, high energy bead (sand) milling or other size reduction processes or mixing means known in the art for effecting particle size reduction in forming a dispersion.

[0026] The organophotoreceptor may include additional layers as well. Such layers are well-known and include, for example, barrier layers, release layers, adhesive layer, and sub-layer. The release layer forms the uppermost layer of the photoconductor element with the barrier layer sandwiched between the release layer and the photoconductive element. The adhesive layer locates and improves the adhesion between the barrier layer and the release layer. The sub-layer is a charge blocking layer and locates between the electrically conductive substrate and the photoconductive element. The sub-layer may also improve the adhesion between the electrically conductive substrate and the photoconductive element.

[0027] Suitable barrier layers include coatings such as crosslinkable siloxanol-colloidal silica coating and hydroxylated silsesquioxane-colloidal silica coating, and organic binders such as polyvinyl alcohol, methyl vinyl ether/maleic anhydride copolymer, casein, polyvinyl pyrrolidone, polyacrylic acid, gelatin, starch, polyurethanes, polyimides, polyesters, polyamides, polyvinyl acetate, polyvinyl chloride, polyvinylidene chloride, polycarbonates, polyvinyl butyral, polyvinyl acetoacetal, polyvinyl formal, polyacrylonitrile, polymethyl methacrylate, polyacrylates, polyvinyl carbazoles, copolymers of monomers used in the above-mentioned polymers, vinyl chloride/vinyl acetate/vinyl alcohol terpolymers, vinyl chloride/vinyl acetate/maleic acid terpolymers, ethylene/vinyl acetate copolymers, vinyl chloride/vinylidene chloride copolymers, cellulose polymers, and mixtures thereof. The above organic binders optionally may contain small inorganic particles such as fumed silica, silica, titania, alumina, zirconia, or a combination thereof. The typical particle size is in the range of 0.001 to 0.5 micrometers, preferably 0.005 micrometers. A preferred barrier layer is a 1:1 mixture of methyl cellulose and methyl vinyl ether/maleic anhydride copolymer with glyoxal as a crosslinker.

[0028] The release layer topcoat may comprise any release layer composition known in the art. Preferably, the release layer is a fluorinated polymer, siloxane polymer, fluorosilicone polymer, silane, polyethylene, polypropylene, polyacrylate, or a combination thereof. More preferably, the release layers are crosslinked silicone polymers.

[0029] Typical adhesive layers include film forming polymers such as polyester, polyvinylbutyral, polyvinylpyrrolidone, polyurethane, polymethyl methacrylate, poly(hydroxy amino ether) and the like. Preferably, the adhesive layer is poly(hydroxy amino ether). If such layers are utilized, they preferably have a dry thickness between about 0.01 micrometer and about 5 micrometers.

[0030] Typical sub-layers include polyvinylbutyral, organosilanes, hydrolyzable silanes, epoxy resins, polyesters, polyamides, polyurethanes, silicones and the like. Preferably, the sub-layer has a dry thickness between about 20 Angstroms and about 2,000 Angstroms.

[0031] The charge transport materials, and organophotoreceptors including these compounds, are suitable for use in an imaging process with either dry or liquid toner development. Liquid toner development is generally preferred because it offers the advantages of providing higher resolution images and requiring lower energy for image fixing compared to dry toners. Examples of useful liquid toners are well-known. They typically include a colorant, a resin

binder, a charge director, and a carrier liquid. A preferred resin to pigment ratio is 2:1 to 10:1, more preferably 4:1 to 8:1. Typically, the colorant, resin, and the charge director form the toner particles.

[0032] The invention will now be described further by way of the following illustrative and non-limiting examples. These examples are to be viewed as being illustrative of specific materials falling within the broader disclosure presented above and are not to be viewed as limiting the broader disclosure.

EXAMPLES

A. Synthesis

(4-n-Butoxycarbonyl-9-fluorenylidene)malononitrile

[0033] To a 1-liter 3-neck round bottom flask, equipped with thermometer, mechanical stirrer and reflux condenser were added 460 g of concentrated sulfuric acid (4.7 moles, analytical grade, commercially obtained from Sigma-Aldrich, Milwaukee, WI) and 100 g of diphenic acid (0.41 mole, commercially obtained from Acros Fisher Scientific Company Inc., Hanover Park, IL). Using heating mantle, the flask was heated to 135-145 °C for 12 minutes, and then cooled to RT. After cooled to RT, the solution was added to a 4 liter Erlenmeyer containing 3 liter of water. The mixture was stirred mechanically and was boiled gently for one hour. A yellow solid was filtered out hot, washed with hot water until the pH of the washing water was neutral, and dried in the air overnight. The yellow solid was fluorenone-4-carboxylic acid (75 g, 80% yield, m.p. 223-224 °C). A ¹H-NMR spectrum of fluorenone-4-carboxylic acid was obtained in d₆-DMSO by a 300 MHz NMR from Bruker Instrument. The peaks were found at δ = 7.39-7.50 (m, 2H); δ = 7.79 - 7.70 (q, 2H); δ = 7.74 -7.85 (d, 1H); δ = 7.88 -8.00 (d, 1H); and δ = 8.18 - 8.30 (d, 1H), where d is doublet, t is triplet, m is multiplet; dd is double doublet, q is quintet.

[0034] To a 2-liter round bottom flask equipped with a mechanical stirrer and a reflux condenser with a Dean Stark apparatus were added 70 g (0.312 mole) of fluorenone-4-carboxylic acid, 480 g (6.5 mole) of n-Butanol (commercially obtained from Fisher Scientific Company Inc., Hanover Park, IL), 1000 ml of Toluene and 4 ml of concentrated sulfuric acid. The solution was refluxed for 5 hours with aggressive agitation and refluxing, during which ~ 6 g of water were collected in the Dean Stark apparatus. The flask was cooled to room temperature. The solvents were evaporated and the residue was added to 4-liter of 3% sodium bicarbonate aqueous solution with agitation. The solid was filtered off, washed with water until the pH of the water was neutral, and dried in the hood overnight. The product was n-butyl fluorenone-4-carboxylate ester (70 g, 80% yield). A ¹H-NMR spectrum of n-butyl fluorenone-4-carboxylate ester was obtained in CDCl₃ by a 300 MHz NMR from Bruker Instrument. The peaks were found at δ = 0.87 -1.09 (t, 3H); δ = 1.42 - 1.70 (m, 2H); δ = 1.75 - 1.88 (q, 2H); δ = 4.26 -4.64 (t, 2H); δ = 7.29 -7.45 (m, 2H); δ = 7.46 -7.58 (m, 1H); δ = 7.60 - 7.68 (dd, 1H); δ = 7.75 - 7.82 (dd, 1H); δ = 7.90 -8.00 (dd, 1H); δ = 8.25 - 8.35 (dd, 1H).

[0035] To a 2-liter, 3-neck round bottom flask equipped with a mechanical stirrer and a reflux condenser were added 70 g (0.25 mole) of n-butyl fluorenone-4-carboxylate ester, 750 ml of absolute methanol, 37 g (0.55 mole) of malononitrile (commercially obtained from Sigma-Aldrich, Milwaukee, WI), 20 drops of piperidine (commercially obtained from Sigma-Aldrich, Milwaukee, WI). The solution was refluxed for 8 hours and the flask was cooled to room temperature. The orange crude product was filtered, washed twice with 70 ml of methanol and once with 150 ml of water, and dried in the hood for overnight. This orange crude product was recrystallized from a mixture of 600 ml of acetone and 300 ml of methanol using activated charcoal. The flask was placed at 0°C for 16 hours. The crystals were filtered and dried in a vacuum oven at 50°C for 6 hours to obtain 60 g of pure (4-n-butoxycarbonyl-9-fluorenylidene) malononitrile. The m.p. was 99-100°C. A ¹H-NMR spectrum of (4-n-butoxycarbonyl-9-fluorenylidene) malononitrile was obtained in CDCl₃ by a 300 MHz NMR from Bruker Instrument. The peaks were found at δ = 0.74 - 1.16 (t, 3H); δ = 1.38 - 1.72 (m, 2H); δ = 1.70 -1.90 (q, 2H); δ = 4.29 - 4.55 (t, 2H); δ = 7.31 - 7.43 (m, 2H); δ = 7.45 - 7.58 (m, 1H); δ = 7.81 - 7.91 (dd, 1H); δ = 8.15 - 8.25 (dd, 1H); δ = 8.42 - 8.52 (dd, 1H); δ = 8.56 -8.66 (dd, 1H).

1,10-Bis(9-fluorenone-4-carboxyl)decane

[0036] 9-Fluorenone-4-carbonyl chloride (2.44 g, 10 mmol) and 1, 10-decanediol (0.87 g, 5 mmol) in the presence of triethylamine (1.01 g, 10 mmol) were refluxed overnight in THF (50 mL). THF was removed in vacuum and the compound was recrystallized using ethyl acetate to give yellow crystals; yield 64%; mp 101-102.1 °C; ¹H-NMR in CDCl₃ (300 MHz) chemical shifts (ppm): 1.34-1.46 (m, 12H), 1.81 (quin, J= 7.2Hz, 4H), 4.41 (t, 8.4Hz, 4H), 7.32-7.37 (m, 4H), 7.50 (td, J=7.8Hz, 2H), 7.70 (d, J=7.2Hz, 2H), 7.82 (dd, J= 1.2Hz, 7.5Hz, 2H), 7.93 (dd, J=0.9Hz, 7.8Hz, 2H), 8.28 (d, J=7.8Hz, 2H). ¹³C-NMR in CDCl₃ (75 MHz) chemical shifts (ppm): 26.14, 28.6, 29.2, 29.4, 65.8, 124.1, 126.2, 127.1, 127.3, 128.6, 129.8, 134.4, 135.1, 135.6, 136.0, 143.2, 144.0, 166.8, 193.0.

1,8-Bis(9-fluorenone-4-carboxyl)octane

[0037] 1,8-Bis(9-fluorenone-4-carboxyl)octane may be prepared according to the following procedure. 9-Fluorenone-4-carboxyl chloride (2.44 g, 10 mmol) and 1,8-octanediol (0.73 g, 5 mmol) in the presence of triethylamine (1.01 g, 10 mmol) are refluxed overnight in THF (50 mL). THF is removed in vacuum and the compound is recrystallized using ethyl acetate to give yellow crystals.

Compound (2)

[0038] A mixture of 4-Methylsulphonylphenylhydrazine hydrochloride (3.5 g, 11.2 mmol), 1,10-bis(9-Fluorenone-4-carboxyl)decane (3.28 g, 5.6 mmol) and sodium acetate (0.92 g, 11.2 mmol) in ethanol (50 mL) was refluxed for 5 h. The resulting mixture was cooled to 20 °C. The precipitate was filtered. The solid was suspended in 100 mL of water and filtered. The filtered cake was washed with ethanol and water to give pure Compound (2) as yellow prisms; yield 45%; m.p. 178-179 °C; ¹H-NMR in CDCl₃ (300MHz) chemical shifts (ppm): 1.33-1.43 (m, 12H), 1.77-1.80 (m, 4H), 3.06 (s, 3H), 3.07 (s, 3H), 4.32-4.38 (m, 2H), 4.40-4.47 (m, 2H), 7.26-7.42 (m, 10H), 7.62-8.05 (m, 10H), 9.04-9.18 (m, 2H)

Compound (3)

[0039] Compound (3) can be prepared by the following procedure. A mixture of 4-Methylsulphonylphenylhydrazine hydrochloride (3.5 g, 11.2 mmol), 1,8-Bis(9-fluorenone-4-carboxyl)octane (3.12 g, 5.6 mmol) and sodium acetate (0.92 g, 11.2 mmol) in ethanol (50 mL) was refluxed for 5 h. The resulting mixture was cooled to 20 °C. The precipitate was filtered. The solid was suspended in 100 mL of water and filtered. The filtered cake was washed with ethanol and water to give pure Compound (3) as yellow prisms.

N-(5-Benzotriazolyl)-N-phenylhydrazine

[0040] N-(5-benzotriazolyl)-N-phenylhydrazine can be prepared according to the procedure described below. To a mixture of phenylhydrazine (97 g, 0.9 mole, commercially available from Aldrich, Milwaukee, WI) and 5-chlorobenzotriazole (15.4 g, 0.1 mole, commercially available from Aldrich, Milwaukee, WI) heated to boiling temperature, sodium is slowly added until there is no more discharge of red coloration. After boiling for some time the mixture is cooled to room temperature. The product is isolated and purified.

Compound (4)

[0041] Compound (4) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and N-(5-benzotriazolyl)-N-phenylhydrazine (29.25 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (4) is isolated and purified.

Compound (5)

[0042] Compound (5) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and N-(5-benzotriazolyl)-N-phenylhydrazine (29.25 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (5) is isolated and purified.

N-Phenyl-N-sulfolan-3-ylhydrazine

[0043] N-Phenyl-N-sulfolan-3-ylhydrazine can be prepared according to the procedure described in Great Britain Patent No. 1,047,525 by Mason, which is incorporated herein by reference. To a mixture of 0.5 mole of butadiene sulfone (commercially available from Aldrich, Milwaukee, WI) and 0.55 mole of phenylhydrazine (commercially available from Aldrich, Milwaukee, WI) was added 0.005 mole 40% aqueous potassium hydroxide solution. The mixture was kept for 2 hours at 60°C whereupon a solid separated. After 10 hours the solid was filtered off to give N-phenyl-N-sulfolan-3-ylhydrazine (53%) having a melting point of 120-121°C (MeOH).

Compound (6)

[0044] A mixture of 1,10-bis(9-fluorenone-4-carboxyl)decane (5.86 g, 10 mmol) and N-phenyl-N-sulfolan-3-ylhydrazine (4.6 g, 20 mmol) was refluxed in ethanol for 5 hours. The mixture was cooled to 0 °C. The compound was filtered

and orange fine crystals of pure Compound (6) were obtained; yield 42% ; m.p. 115.9-116.1 °C; ¹H-NMR in CDCl₃ (300MHz) chemical shifts (ppm): 1.19-1.42 (m, 12H), 1.76-1.81 (m, 4H) , 2.57-2.71 (m, 4H), 3.08-3.17(m, 2H), 3.24-3.33 (m, 2H), 3.48-3.59 (m, 2H), 3.74-3.85 (m, 2H), 4.38-4.42 (m, 4H), 4.77 (t, J=7.0Hz, 2H), 6.94-7.07 (m, 8H) , 7.2-7.35 (m, 8H), 7.37-7.42 (m, 2H), 7.50 (d, J=7.8Hz, 1H), 7.70 (dd, J=7.8Hz, 16.8Hz, 1H), 7.84 (d, J=7.5Hz, 1H), 8.03-8.12 (m, 1H), 8.22 (t, 7.8Hz, 2H).

Compound (7)

[0045] A mixture of 1,8-Bis(9-fluorenone-4-carboxyl)octane (5.58 g, 10 mmol) and N-phenyl-N-sulfolan-3-ylhydrazine (4.6 g, 20 mmol) was refluxed in ethanol for 5 hours. The mixture was cooled 0 °C. The compound was filtered and orange fine crystals of pure Compound (7) were obtained.

5-Methyl-1-Phenyl-3-(1-phenylhydrazino)pyrazole

[0046] 5-Methyl-1-phenyl-3-(1-phenylhydrazino)pyrazole can be prepared according to the procedure described in J. Chem. Soc. C (1971), (12), 2314-17 by Boyd et al., which is incorporated herein by reference.

Compound (8)

[0047] Compound (8) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and 5-methyl-1-phenyl-3-(1-phenylhydrazino)pyrazole (34.32 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (8) is isolated and purified.

Compound (9)

[0048] Compound (9) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and 5-methyl-1-phenyl-3-(1-phenylhydrazino)pyrazole (34.32 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (9) is isolated and purified.

N-(4-Stilbenyl)-N-phenylhydrazine

[0049] N-(4-Stilbenyl)-N-phenylhydrazine can be prepared according to the procedure described in Zh. Org. Khim. (1967), 3(9), 1605-3 by Matevosyan et al., which is incorporated herein by reference. To a mixture of phenylhydrazine (97 g, 0.9 mole, commercially available from Aldrich, Milwaukee, WI) and p-chlorostilbene (21.4 g, 0.1 mole, commercially available from Spectrum Quality Products, Inc., Gardena, CA; Web: www.spectrumchemical.com) heated to boiling temperature, sodium was slowly added until there was no more discharge of red coloration. After boiling for some time the mixture was dissolved in 1750 ml of ethanol and cooled to -15 °C. The precipitated product was recrystallized to give 28% of N-(4-stilbenyl)-N-phenylhydrazine.

Compound (10)

[0050] Compound (10) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and N-(4-stilbenyl)-N-phenylhydrazine (37.18 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (10) is isolated and purified.

Compound (11)

[0051] Compound (11) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and N-(4-stilbenyl)-N-phenylhydrazine (37.18 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (11) is isolated and purified.

1-Phenyl-1-(1-benzyl-1H-tetrazol-5-yl)hydrazine

[0052] 1-Phenyl-1-(1-benzyl-1H-tetrazol-5-yl)hydrazine can be prepared according to the procedure described in Tetrahedron (1983), 39(15), 2599-608 by Atherton et al., which is incorporated herein by reference.

Compound (12)

[0053] Compound (12) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and 1-phenyl-1-(1-benzyl-1H-tetrazol-5-yl)hydrazine (34.58 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (12) is isolated and purified.

Compound (13)

[0054] Compound (13) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and 1-phenyl-1-(1-benzyl-1H-tetrazol-5-yl)hydrazine (34.58 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (13) is isolated and purified.

1,1-Dinaphthylhydrazine

[0055] 1,1-Dinaphthylhydrazine can be prepared according to the procedure described in Staschkow, L.I.; Matevosyan, R.O. Journal of the General Chemistry (1964) 34, 136, which is incorporated herein by reference. A suspension of 0.07 mole of the naphthyl nitrosamine in 750 ml of ether is cooled to 5-8 °C and treated with 150 g of zinc dust. Acetic acid (70 ml) is then added drop wise with stirring. To complete the reaction, 40g of zinc dust is added. The reaction mixture is heated and filtered from the sludge. The mother liquor is washed with 10% sodium carbonate solution and dried with solid KOH. The ether is distilled off to give the crystalline hydrazine, which is crystallized from ethanol or butanol.

Compound (14)

[0056] Compound (14) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and dinaphthylhydrazine (37.18 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (14) is isolated and purified.

Compound (15)

[0057] Compound (15) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and dinaphthylhydrazine (37.18 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (15) is isolated and purified.

N-4-[(9H-Fluoren-9-ylidene)benzyl]-N-phenylhydrazine

[0058] N-4-[(9H-fluoren-9-ylidene)benzyl]-N-phenylhydrazine can be prepared according to the procedure similar to that described in Zh. Org. Khim. (1967), 3(9), 1605-3 by Matevosyan et al., which is incorporated herein by reference. To a mixture of phenylhydrazine (97 g, 0.9 mole, commercially available from Aldrich, Milwaukee, WI) and p-9-(4-chlorobenzylidene)fluorene (28.9 g, 0.1 mole, commercially available from Aldrich, Milwaukee, WI) heated to boiling temperature, sodium was slowly added until there was no more discharge of red coloration. After boiling for some time the mixture was dissolved in 1750 ml of ethanol and cooled to -15 °C. The precipitated product was recrystallized to give N-4-[(9H-fluoren-9-ylidene)benzyl]-N-phenylhydrazine.

Compound (16)

[0059] Compound (16) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and N-4-[(9H-fluoren-9-ylidene)benzyl]-N-phenylhydrazine (46.8 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (16) is isolated and purified.

Compound (17)

[0060] Compound (17) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and N-4-[(9H-fluoren-9-ylidene)benzyl]-N-phenylhydrazine (46.8 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (17) is isolated and purified.

N-Pyrrol-2-yl-N-phenylhydrazine

[0061] N-Pyrrol-2-yl-N-phenylhydrazine can be prepared according to the procedure described in Japanese Patent No. 05148210 by Myamoto, which is incorporated herein by reference.

Compound (18)

[0062] Compound (18) can be prepared by the following procedure. 1,10-Bis(9-fluorenone-4-carboxyl)decane (37.5 g, 64 mmol) and N-pyrrol-2-yl-N-phenylhydrazine (22.49 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (18) is isolated and purified.

Compound (19)

[0063] Compound (19) can be prepared by the following procedure. 1,8-Bis(9-fluorenone-4-carboxyl)octane (35.71 g, 64 mmol) and N-pyrrol-2-yl-N-phenylhydrazine (22.49 g, 130 mmol, 2 equiv.) are combined in tetrahydrofuran (600 mL) and heated at reflux with stirring for 2 hours. Compound (19) is isolated and purified.

Example 1

[0064] Example 1 was a single layer organophotoreceptor that was ring coated onto a 30 mm x 250 mm, anodized aluminum drum core at a setting to produce a nominal thickness of 13 microns. The coating solution for the single layer organophotoreceptor was prepared by combining 1.87 g of MPCT-10 (a charge transfer material, commercially obtained from Mitsubishi Paper Mills, Tokyo, Japan), 0.54 g of Compound 2, and 9.37 g of tetrahydrofuran and shaken until the components dissolved. Added to this mixture was 7.4 g of a 14wt% polyvinyl butyral resin (BX-1, commercially obtained from Sekisui Chemical Co. Ltd., Japan) in tetrahydrofuran pre-mix solution and 0.83 g of a CGM mill-base containing 18.5wt% of titanyl oxyphthalocyanine plus polyvinyl butyral resin at a ratio of 2.3:1 (BX-5, commercially obtained from Sekisui Chemical Co. Ltd., Japan) in tetrahydrofuran. Tinuvin 292 (0.018 g T-292 in 5.68 g tetrahydrofuran) light stabilizer was added to the coating solution.

[0065] The CGM mill-base was obtained by milling 112.7 g of titanyl oxyphthalocyanine (commercially obtained from H.W. Sands Corp., Jupiter, FL) with 49 g of the polyvinyl butyral resin (BX-5) in 651 g of MEK on a horizontal sand mill (model LMC12 DCMS, commercially obtained from Netzsch Incorporated, Exton, PA) with 1-micron zirconium beads using recycle mode for 4 hours.

[0066] After mixing the solution on a mechanical shaker for ~ 1 hour, the single layer coating solution was coated onto drum core substrate described above and dried in a convection oven at 100°C for 15-30 minutes.

Example 2

[0067] Example 2 single layer organophotoreceptor coating solution was prepared by combining 1.87 g of MPCT-10 (a charge transfer material, commercially obtained from Mitsubishi Paper Mills, Tokyo, Japan), 0.54 g of Compound 6 and 9.37 g of tetrahydrofuran and shaken until the components dissolved. Added to this mixture was 7.4 g of a 14wt% polyvinyl butyral resin (BX-1, commercially obtained from Sekisui Chemical Co. Ltd., Japan) in tetrahydrofuran pre-mix solution and 0.83 g of a CGM mill-base containing 18.5wt% of titanyl oxyphthalocyanine plus polyvinyl butyral resin at a ratio of 2.3:1 (BX-5, commercially obtained from Sekisui Chemical Co. Ltd., Japan) in tetrahydrofuran. Tinuvin 292 (0.018 g T-292 in 5.68 g tetrahydrofuran) light stabilizer was added to the coating solution.

[0068] The CGM mill-base was prepared as described in Example 1.

[0069] After mixing the solution on a mechanical shaker for ~ 1 hour, the single layer coating solution was coated onto the drum substrate described in Example 1 using a ring coater and dried in a convection oven at 100°C for 15-30 minutes.

Comparative Example A

[0070] Comparative Example A single layer organophotoreceptor coating solution was prepared by combining 1.87 g of MPCT-10 (a charge transfer material, commercially obtained from Mitsubishi Paper Mills, Tokyo, Japan), 0.54 g of a (4-n-butoxycarbonyl-9-fluorenylidene)malononitrile, and 9.37 g of tetrahydrofuran and shaken until the components dissolved. Added to this mixture was 7.4 g of a 14wt% polyvinyl butyral resin (BX-1, commercially obtained from Sekisui Chemical Co. Ltd., Japan) in tetrahydrofuran pre-mix solution and 0.83 g of a CGM mill-base containing 18.5wt% of titanyl oxyphthalocyanine plus polyvinyl butyral resin at a ratio of 2.3:1 (BX-5, commercially obtained from Sekisui Chemical Co. Ltd., Japan) in tetrahydrofuran. Tinuvin 292 (0.018 g T-292 in 5.68 g tetrahydrofuran) light stabilizer was

added to the coating solution.

[0071] The CGM mill-base was prepared as described in Example 1.

After mixing the solution on a mechanical shaker for ~ 1 hour, the single layer coating solution was coated onto the substrate described in Example 1 using a ring coater and dried in a convection oven at 100°C for 15-30 minutes.

Electrostatic Testing

[0072] Electrostatic cycling performance of the charge transfer compounds of this invention is determined using in-house designed and developed test bed that is capable of testing the ring coated 30 mm drums. Electrostatic evaluation on the 30 mm drum test bed is designed to accelerate electrostatic fatigue during extended cycling by increasing the charge-discharge cycling frequency and decreasing the recovery time as compared to larger diameter drum formats. In this electrostatic cycling tester, the drum rotated at a rate of 12.7 cm / sec (5 ips) and the location of each station in the tester (distance and elapsed time per cycle) is given as:

[0073] Electrostatic test stations around the 30 mm drum at 12.7 cm / sec.

Station	Degrees	Total Distance, cm	Total Time, sec
Erase Bar Center	0°	Initial, 0 cm	Initial, 0 s
Scorotron Charger	87.3°	2.28	0.18
Laser Strike	147.7°	3.86	0.304
Probe #1	173.2°	4.53	0.357
Probe #2	245.9°	6.43	0.506
Erase Bar Center	360°	9.42	0.742

[0074] From the above table, the first electrostatic probe (Trek 344 electrostatic meter) is located 0.05 s after the laser strike station and 0.18 s after the scorotron while the second probe (Trek 344 electrostatic meter) is located 0.15 s from the first probe and 0.33 s from the scorotron. All measurements were performed at 20°C and 30% relative humidity.

[0075] Electrostatic measurements were obtained as a compilation of several tests. The first three diagnostic tests (prodstart, VlogE initial, dark decay initial) are designed to evaluate the electrostatic cycling of a new, fresh sample and the last three, identical diagnostic test (prodend, VlogE final, dark decay final) are run after cycling of the sample (longrun). The laser is operated at 780nm, 600dpi, 50 um spot size, 60 nanoseconds / pixel expose time, 1,800 lines per second scan speed, and a 100% duty cycle. The duty cycle is the percent exposure of the pixel clock period, i.e., the laser is on for the full 60 nanoseconds per pixel at a 100% duty cycle.

Electrostatic Test Suite:

[0076]

1) PRODTEST: The erase bar was turned on during this diagnostic test and the sample recharged at the beginning of each cycle (except where indicated as charger off). The test sequence was as follows. The sample was completely charged for three complete drum revolutions (laser off); discharged with the laser (780nm, 600dpi, 50 um spot size, expose 60 nanoseconds / pixel, run at a scan speed of 1,800 lines per second, and use a 100% duty cycle) on the fourth cycle; completely charged for the next three cycles (laser off); discharged with only the erase lamp @ 720nm on the eighth cycle (corona and laser off); and, finally, completely charged for the last three cycles (laser off).

2) VLOGE: This test measures the photoinduced discharge of the photoconductor to various laser intensity levels by monitoring the discharge voltage of the belt as a function of the laser power (exposure duration of 50 ns). The complete sample was charged and discharged at incremental laser power levels per each drum revolution. A semi-logarithmic plot was generated (voltage verses log E) to identify the sample's sensitivity and operational power settings. The functional photosensitivity (Sens.) was determined from this diagnostic test.

3) DARK DECAY: This test measures the loss of charge acceptance in the dark with time without laser or erase illumination and can be used as an indicator of i) the injection of residual holes from the charge generation layer to the charge transport layer, ii) the thermal liberation of trapped charges, and iii) the injection of charge from the surface or aluminum ground plane. After the belt has been completely charged, it was stopped and the probes

measured the surface voltage over a period of 90 seconds. The decay in the initial voltage was plotted versus time.

4) LONGRUN: The belt was electrostatically cycled for 100 to 4,000 drum revolutions according to the following sequence per each belt-drum revolution. The belt was charged by the corona, the laser was cycled on and off (80-100° sections) to discharge a portion of the belt and, finally, the erase lamp discharged the whole belt in preparation for the next cycle. The laser was cycled so that the first section of the belt was never exposed, the second section was always exposed, the third section was never exposed, and the final section was always exposed. This pattern was repeated for a total of 100 to 4,000 drum revolutions and the data was recorded periodically, after every "total drum revolutions"/twenty cycles.

5) After the last cycle (long run test), the PRODTST, VLOGE, DARK DECAY diagnostic tests were run again.

[0077] The following Table shows the results from the prodstart and prodend diagnostic tests. The values for the charge acceptance voltage (Vacc, probe #1 average voltage obtained from the third cycle), discharge voltage (Vdis, probe #1 average voltage obtained from the fourth cycle), functional dark decay voltage (Vdd, average voltage difference between probes 1 & 2 obtained from the third cycle), and the residual voltage (Vres, probe 1, average voltage obtained from the eighth cycle) are reported for the initial and final cycles.

Table 1 . The Electrostatic Test Results after 500 Cycles.

Sample	Prodstart					Prodstop				
	Vacc	Vdis	Sens.	Vdd	Vres	Vacc	Vdis	Sens.	Vdd	Vres
Comparative Example A	601	40	419	51	51	586	39	397	39	9
Example 1	863	150	236	20	61	625	145	236	23	70
Example 2	740	128	222	28	59	523	103	222	29	58

[0078] In the above table, the radiation sensitivity (Sensitivity at 780nm in m2/J) of the xerographic process was determined from the information obtained during the VLOGE diagnostic run by calculating the reciprocal of the product of the laser power required to discharge the photoreceptor to ½ of its initial potential, the exposure duration, and 1/spot size.

[0079] A number of embodiments of the invention have been described. Nevertheless, it will be understood that various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.

[0080] Attention is directed to all papers and documents which are filed concurrently with or previous to this specification in connection with this application and which are open to public inspection with this specification, and the contents of all such papers and documents are incorporated herein by reference.

[0081] All of the features disclosed in this specification (including any accompanying claims, abstract and drawings), and/or all of the steps of any method or process so disclosed, may be combined in any combination, except combinations where at least some of such features and/or steps are mutually exclusive.

[0082] Each feature disclosed in this specification (including any accompanying claims, abstract and drawings) may be replaced by alternative features serving the same, equivalent or similar purpose, unless expressly stated otherwise. Thus, unless expressly stated otherwise, each feature disclosed is one example only of a generic series of equivalent or similar features.

[0083] The invention is not restricted to the details of the foregoing embodiment(s). The invention extends to any novel one, or any novel combination, of the features disclosed in this specification (including any accompanying claims, abstract and drawings), or to any novel one, or any novel combination, of the steps of any method or process so disclosed.


Claims

1. An organophotoreceptor comprising:

- (a) at least one charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolyhydrazone groups, d) at least two fluorenone sulfolanylhydrazone groups, e) at least two fluorenone pyrazolyhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolyhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups;
 (b) a charge generating compound; and
 (c) an electrically conductive substrate.

2. An organophotoreceptor comprising:

- (a) a charge transport material having a formula selected from the group consisting of:



A) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is an alkylsulfonylphenyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

B) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a pyrrolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

C) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a benzotriazolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

D) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a sulfolanyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

E) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a pyrazolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

F) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a tetrazolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

G) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a stilbenyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom,

a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ -group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

H) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a (9H-fluoren-9-ylidene)benzyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ -group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group; and

I) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a naphthyl group or one of its derivatives;


X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ -group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

(b) a charge generating compound; and

(c) an electrically conductive substrate.

3. The organophotoreceptor of claim 1 wherein the at least one charge transport material comprises a compound of the formula:



where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is an alkylsulfonylphenyl group, pyrrolyl group, benzotriazolyl group, sulfolanyl group, pyrazolyl group, tetrazolyl group, stilbenyl group, (9H-fluoren-9-ylidene)benzyl group, naphthyl group, or one of their derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are,

independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

4. An organophotoreceptor according to claim 2 wherein said organophotoreceptor further comprises a second charge transport material.

5. An organophotoreceptor according to claim 2 comprising:


- (a) a charge transport layer comprising said charge transport material and a polymeric binder;
- (b) a charge generating layer comprising said charge generating compound and a polymeric binder; and
- (c) said electrically conductive substrate.

6. An electrophotographic imaging apparatus comprising:

- (a) a plurality of support rollers; and
- (b) an organophotoreceptor in the form of a flexible belt threaded around said support rollers, said organophotoreceptor comprising:

- (i) at least one charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolyhydrazone groups, d) at least two fluorenone sulfanylhydrazone groups, e) at least two fluorenone pyrazolyhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolyhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups;
- (ii) a charge generating compound; and
- (iii) an electrically conductive substrate.

7. The electrophotographic imaging apparatus of claim 6 wherein the charge transport material is from the group consisting of:



A) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is an alkylsulfonylphenyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a hetero-

cyclic group, or a CR₇ group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

B) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a pyrrolyl group or one of its derivatives;

X is a linking group having the formula -(CH₂)_m-, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a -NR₃ group, a CHR₄ group, or a CR₅R₆ group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear -(CH₂)_p- group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR₇ group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

C) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a benzotriazolyl group or one of its derivatives;

X is a linking group having the formula -(CH₂)_m-, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a -NR₃ group, a CHR₄ group, or a CR₅R₆ group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear -(CH₂)_p- group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR₇ group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

D) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a sulfolanyl group or one of its derivatives;

X is a linking group having the formula -(CH₂)_m-, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a -NR₃ group, a CHR₄ group, or a CR₅R₆ group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear -(CH₂)_p- group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR₇ group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

E) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a pyrazolyl group or one of its derivatives;

X is a linking group having the formula -(CH₂)_m-, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a -NR₃ group, a CHR₄ group, or a CR₅R₆ group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear -(CH₂)_p- group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR₇ group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

F) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a tetrazolyl group or one of its derivatives;

X is a linking group having the formula -(CH₂)_m-, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a -NR₃ group, a CHR₄ group, or a CR₅R₆ group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

G) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a stilbenyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

H) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a (9H-fluoren-9-ylidene)benzyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group; and

I) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a naphthyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

8. An electrophotographic imaging process comprising:

(a) applying an electrical charge to a surface of an organophotoreceptor comprising:

(i) at least one charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazone groups, b) at least two fluorenone pyrrolylhydrazone groups, c) at least two fluorenone benzotriazolylhydrazone groups, d) at least two fluorenone sulfolanylhydrazone groups, e) at least two fluorenone pyrazolylhydrazone groups, f) at least two fluorenone naphthylhydrazone groups, g) at least two fluorenone tetrazolylhydrazone groups, h) at least two fluorenone stilbenylhydrazone groups, and i) at least two fluorenone (9H-fluoren-9-ylidene)benzylhydrazone groups;

(ii) a charge generating compound; and


(iii) an electrically conductive substrate;

(b) imagewise exposing said surface of said organophotoreceptor to radiation to dissipate charge in selected areas and thereby form a pattern of charged and uncharged areas on said surface;

(c) contacting said surface with a liquid toner comprising a dispersion of colorant particles in an organic liquid to create a toned image; and

(d) transferring said toned image to a substrate.

9. The imaging process of claim 8 wherein the charge transport material comprises a compound of the formula



wherein the formula is defined by a combination of groups selected from the group consisting of:

A) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is an alkylsulfonylphenyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

B) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a pyrrolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

C) where n is an integer between 2 and 6, inclusive;

R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is a benzotriazolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group;

D) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a sulfolanyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

E) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a pyrazolyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

F) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a tetrazolyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

G) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a stilbenyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

H) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a (9H-fluoren-9-ylidene)benzyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group; and

l) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;


R_2 is a naphthyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

10. A charge transport material comprising a fluorenone hydrazone having a combination of groups thereon, the combination of groups being selected from the group consisting of a) at least two fluorenone alkylsulfonylphenylhydrazones, b) at least two fluorenone pyrrolylhydrazones, c) at least two fluorenone benzotriazolyldrazones, d) at least two fluorenone sulfonanyldrazones, e) at least two fluorenone pyrazolyldrazones, f) at least two fluorenone naphthylhydrazones, g) at least two fluorenone tetrazolyldrazones, h) at least two fluorenone stilbenylhydrazones, and i) at least two fluorenone (9H-fluoren-9-ylidene) benzyldrazones.

11. The charge transport material of claim 10 wherein the charge transport material comprises a compound of the formula:



wherein the formula is defined by definitions selected from the group consisting of:

A) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is an alkylsulfonylphenyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

B) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a pyrrolyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 ,

R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group
 where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a hetero-
 cyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

C) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a benzotriazolyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and
 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl
 group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 ,
 R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group
 where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a hetero-
 cyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

D) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a sulfolanlyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and
 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl
 group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 ,
 and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group
 where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a hetero-
 cyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

E) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a pyrazolyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and
 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl
 group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 ,
 R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group
 where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a hetero-
 cyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

F) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a tetrazolyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and
 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl
 group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 ,
 R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and
 Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group
 where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a hetero-
 cyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

G) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;
 R_2 is a stilbenyl group or one of its derivatives;
 X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and

20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group;

H) where n is an integer between 2 and 6, inclusive;

R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a (9H-fluoren-9-ylidene)benzyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; or

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group; and

I) where n is an integer between 2 and 6, inclusive;


R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is a naphthyl group or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

12. The organophotoreceptor of claim 2 wherein the charge transport material has the formula



where n is an integer between 2 and 6, inclusive;


R_1 is hydrogen, an alkyl group, or an aryl group;

R_2 is an alkylsulfonylphenyl group, pyrrolyl group, benzotriazolyl group, sulfolanyl group, pyrazolyl group, tetrazolyl group, stilbenyl group, (9H-fluoren-9-ylidene)benzyl group, naphthyl group, or one of their derivatives;

X is a linking group having the formula $-(CH_2)_m$ -, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p$ - group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxy group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

13. The electrophotographic imaging apparatus of claim 7 wherein the charge transport material has the formula



where n is an integer between 2 and 6, inclusive;


R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is an alkylsulfonylphenyl group, pyrrolyl group, benzotriazolyl group, sulfolanyl group, pyrazolyl group, tetrazolyl group, stilbenyl group, (9H-fluoren-9-ylidene)benzyl group, naphthyl group, or one of their derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group.

14. The process of claim 9 wherein the charge transport material has the formula



where n is an integer between 2 and 6, inclusive;


R₁ is hydrogen, an alkyl group, or an aryl group;

R₂ is an alkylsulfonylphenyl group, pyrrolyl group, benzotriazolyl group, sulfolanyl group, pyrazolyl group, tetrazolyl group, stilbenyl group, (9H-fluoren-9-ylidene)benzyl group, naphthyl group, or one of their derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R₃, R₄, R₅, and R₆ are, independently, H, an alkyl group, or an aryl group; and


Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R₇ is hydrogen atom, an alkyl group, or an aryl group.

15. The organophotoreceptor of claim 2 wherein the charge transport material has the formula




where m is an integer between 2 and 20.

16. The organophotoreceptor of claim 2 wherein the charge transport material has the formula



where m is an integer between 2 and 20.

17. The organophotoreceptor of claim 2 wherein the charge transport material has the formula



where n is an integer between 2 and 6, inclusive;


R_1 and R_2 are, independently, naphthyl or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and

20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and


Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

18. The charge transport material of claim 10 wherein the charge transport material has the formula




where m is an integer between 2 and 20.

19. The charge transport material of claim 10 wherein the charge transport material has the formula



where m is an integer between 2 and 20.

20. The charge transport material of claim 10 wherein the charge transport material has the formula




15 where n is an integer between 2 and 6, inclusive;




R_1 and R_2 are, independently, naphthyl or one of its derivatives;

X is a linking group having the formula $-(CH_2)_m-$, branched or linear, where m is an integer between 0 and 20, inclusive, and one or more of the methylene groups is optionally replaced by oxygen atom, a carbonyl group, urethane, urea, an ester group, a $-NR_3$ group, a CHR_4 group, or a CR_5R_6 group where R_3 , R_4 , R_5 , and R_6 are, independently, H, an alkyl group, or an aryl group; and

Y is a bond, carbon atom, nitrogen atom, oxygen atom, sulfur atom, a branched or linear $-(CH_2)_p-$ group where p is an integer between 0 and 10, an aryl group, a cycloalkyl group, a cyclosiloxyl group, a heterocyclic group, or a CR_7 group where R_7 is hydrogen atom, an alkyl group, or an aryl group.

25 21. The organophotoreceptor of claim 1 wherein the charge transport material is represented by one of the formulae selected from the group consisting of:





5

10

15

20

25

30




35

40

45

50

55



5

10

15

20

25

30




35




40


45

50

55




















22. The electrophotographic imaging process of claim 8 wherein the charge transport material is represented by one




of the formulae selected from the group consisting of:


























23. The charge transport material of claim 10 wherein the charge transport material is represented by one of the formulae selected from the group consisting of:












5

10

15




20

25

30

35





40

45

50

55







European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 03 25 0767

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	DATABASE WPI Section Ch, Week 198851 Derwent Publications Ltd., London, GB; Class E06, AN 1988-363223 XP002240965 & JP 63 271452 A (MITA), 9 November 1988 (1988-11-09) * abstract *	1,10	G03G5/06 C07D333/48 C07C251/84 C07C317/34 C07D249/18 C07D231/38 C07D257/06 C07D207/34
A	DATABASE WPI Section Ch, Week 199532 Derwent Publications Ltd., London, GB; Class E08, AN 1995-243615 XP002240966 & JP 07 149704 A (MITA), 13 June 1995 (1995-06-13) * abstract *	1,10	
A	DATABASE WPI Section Ch, Week 199508 Derwent Publications Ltd., London, GB; Class E10, AN 1995-054934 XP002240967 & JP 06 332203 A (CANON), 2 December 1994 (1994-12-02) * abstract *	1,10	TECHNICAL FIELDS SEARCHED (Int.Cl.7) G03G C07D C07C
A	US 4 415 640 A (S GOTO) 15 November 1983 (1983-11-15) * claim 1 *	1,10	
A	US 6 004 708 A (M T BELLINO) 21 December 1999 (1999-12-21) * claim 1 *	1,10	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13 May 2003	Examiner Vanhecke, H
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 03 25 0767

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-05-2003

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
JP 63271452	A	09-11-1988	JP	2561924 B2	11-12-1996
JP 7149704	A	13-06-1995	NONE		
JP 6332203	A	02-12-1994	JP	3197117 B2	13-08-2001
US 4415640	A	15-11-1983	JP	1331328 C	14-08-1986
			JP	57138644 A	27-08-1982
			JP	60058469 B	20-12-1985
US 6004708	A	21-12-1999	AU	2963700 A	02-11-2000
			CN	1351722 T	29-05-2002
			EP	1171805 A1	16-01-2002
			JP	2002542515 T	10-12-2002
			WO	0063748 A1	26-10-2000

EPO FORM P4459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.